

ONSAGER-TYPE CONJECTURE AND RENORMALIZED SOLUTIONS FOR THE RELATIVISTIC VLASOV-MAXWELL SYSTEM

BY

CLAUDE BARDOS (*Laboratoire J.-L. Lions, Université Paris Diderot, UFR de Mathématiques Bâtiment Sophie Germain, 75205 Paris Cedex 13, France*),

NICOLAS BESSE (*Laboratoire J.-L. Lagrange, UMR CNRS/OCA/UCA 7293, Université Côte d'Azur, Observatoire de la Côte d'Azur, Bd de l'observatoire CS 34229, 06300 Nice, Cedex 4, France*),

AND

TOAN T. NGUYEN (*Department of Mathematics, Pennsylvania State University, State College, Pennsylvania, 16803*)

This paper is dedicated to Walter Strauss on the occasion of his 80th birthday, as a token of friendship and admiration, in particular, for his contribution to the mathematical theory of Vlasov-Maxwell systems.

Abstract. In this paper we give a proof of an Onsager-type conjecture on conservation of energy and entropies of weak solutions to the relativistic Vlasov–Maxwell equations. As concerns the regularity of weak solutions, say in Sobolev spaces $W^{\alpha,p}$, we determine Onsager-type exponents α that guarantee the conservation of all entropies. In particular, the Onsager exponent α is smaller than $\alpha = 1/3$ established for fluid models. Entropies conservation is equivalent to the renormalization property, which has been introduced by DiPerna–Lions for studying well-posedness of passive transport equations and collisionless kinetic equations. For smooth solutions, renormalization property or entropies conservation are simply the consequence of the chain rule. For weak solutions the use of the chain rule is not always justified. Then arises the question about the minimal regularity needed for weak solutions to guarantee such properties. In the DiPerna–Lions and Bouchut–Ambrosio theories, renormalization property holds under sufficient conditions in terms of the regularity of the advection field, which are roughly speaking an entire derivative in some Lebesgue spaces (DiPerna–Lions) or an entire derivative in the

Received March 9, 2019, and, in revised form, May 27, 2019.

2010 *Mathematics Subject Classification*. Primary 35Q60, 35Q61, 35Q83.

Key words and phrases. Relativistic Vlasov–Maxwell system, Onsager’s conjecture, entropies conservation, renormalization property, energy conservation.

Email address: claude.bardos@gmail.com

Email address: nicolas.besse@oca.eu

Email address: nguyen@math.psu.edu

©2019 Brown University

space of measures with finite total variation (Bouchut–Ambrosio). In return there is no smoothness requirement for the advected density, except some natural a priori bounds. Here we show that the renormalization property holds for an electromagnetic field with only a fractional space derivative in some Lebesgue spaces. To compensate for this loss of derivative for the electromagnetic field, the distribution function requires an additional smoothness, typically fractional Sobolev differentiability in phase-space. As concerns the conservation of total energy, if the macroscopic kinetic energy is in L^2 , then total energy is preserved.

1. Introduction. The dimensionless relativistic Vlasov–Maxwell system reads,

$$\partial_t f + v \cdot \nabla_x f + (E + v \times B) \cdot \nabla_\xi f = 0, \quad (1)$$

$$\partial_t E - \nabla \times B = -j, \quad \partial_t B + \nabla \times E = 0, \quad (2)$$

$$\nabla \cdot E = \rho, \quad \nabla \cdot B = 0, \quad (3)$$

where $t \in \mathbb{R}$, $x \in \mathbb{R}^3$, $\xi \in \mathbb{R}^3$, and $v = \xi / \sqrt{1 + |\xi|^2}$ represent time, position, momentum, and velocity of particles, respectively. The distribution function of particles $f = f(t, x, \xi)$ satisfies the Vlasov equation (1) with acceleration given by the Lorentz force $F_L = E + v \times B$, while the electromagnetic field $E = E(t, x)$ and $B = B(t, x)$ satisfies Maxwell's equations (2)-(3). The coupling between the Vlasov equation and Maxwell's equations occurs through the source terms of Maxwell's equations, which are the charge density $\rho = \rho(t, x)$ and the current density $j = j(t, x)$. These densities are defined as the first v -moments of the phase-space density of particles f , namely,

$$\rho(t, x) = \int_{\mathbb{R}^3} f(t, x, \xi) d\xi, \quad j(t, x) = \int_{\mathbb{R}^3} v f(t, x, \xi) d\xi. \quad (4)$$

The initial value problem associated to the system (1)-(4) requires initial conditions given by,

$$f(0, x, \xi) = f_0(x, \xi) \geq 0, \quad (5)$$

$$E(0, x) = E_0(x), \quad B(0, x) = B_0(x), \quad \nabla \cdot E_0 = \rho_0 = \int_{\mathbb{R}^3} f_0 d\xi, \quad \nabla \cdot B_0 = 0. \quad (6)$$

In addition for the well-posedness of Maxwell's equations (2)-(3), the densities of charge ρ and current j must satisfy a compatibility condition given by the charge conservation law,

$$\partial_t \rho + \nabla \cdot j = 0. \quad (7)$$

This continuity equation is automatically satisfied if the Vlasov equation (1) is satisfied since it can be recovered by integration in the momentum variable of the Vlasov equation. Let us note that Maxwell–Gauss equations (3) are satisfied at any time if they are satisfied initially. Indeed, it is a consequence of time integration of the divergence of the Maxwell–Faraday–Ampère equations (2), in combination with the continuity equation (7) and initial conditions (6).

The Vlasov equation (1) has, at least formally, infinitely many invariants. Indeed, let $\mathcal{H} : \mathbb{R} \rightarrow \mathbb{R}$ be any smooth function. Multiplying (1) with $\mathcal{H}'(f)$ and applying the chain rule, we then obtain,

$$\partial_t \mathcal{H}(f) + v \cdot \nabla_x \mathcal{H}(f) + (E + v \times B) \cdot \nabla_\xi \mathcal{H}(f) = 0. \quad (8)$$

A solution f to (1) in the sense of distributions is said to be a renormalized solution if for any smooth nonlinear function \mathcal{H} , f also solves (8) in the sense of distributions. We say that the field (v, F_L) satisfies the renormalization property if any solution f to (1) in the sense of distributions is a renormalized solution. The renormalization technique appeared in the well-posedness of passive advection equations and ODEs [37], in the analysis of the Boltzmann equation [38], in the theory of weak solutions of the compressible Navier-Stokes equations [57], and in the theory of weak solutions of collisionless kinetic equations such as the Vlasov–Poisson system [34, 35]. The groundbreaking work [37] has highlighted the fundamental link between renormalized solutions to the passive transport equation,

$$\partial_t u + b \cdot \nabla u = 0, \quad u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}, \quad b : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d, \quad (9)$$

and the well-posedness theory for the associated ODE,

$$\partial_t X(t, x) = b(t, X(t, x)), \quad t \in [0, T], \quad X(0, x) = x \in \mathbb{R}^d, \quad (10)$$

where b is a nonsmooth vector field. Similarly to entropy conditions for hyperbolic conservation laws, renormalization property provides additional stability under weak convergence. Indeed, renormalized solutions come with a comparison principle, which allows us to show uniqueness of renormalized solutions and some stability results for sequences of solutions. In return, uniqueness at the PDE level (9) implies uniqueness at the ODE level (10). It was first shown in [37] that the renormalization property holds provided $b \in L_t^1 W_x^{1,p}$ with $p \geq 1$, plus a bounded divergence and a global space growth estimate on b (see also [55] for the case $W_{\text{loc}}^{1,1}$). Moreover, there is no additional regularity assumption for u except its boundedness or some L^p -bounds. This result was extended to $b \in L_t^1 BV_x$ with $\nabla \cdot b \in L_{t,x}^1$, first in [23] for the Vlasov equation (see also [53] for a related result), and then in [9] for the general case (see also [28]). Very recently, in [11] the authors developed a local version of the DiPerna–Lions’ theory under no global assumptions on the growth estimate of b . We refer the reader to [10] for a recent survey.

For the Vlasov–Poisson system when f is merely L^1 , the product Ef does not belong to L_{loc}^1 . Therefore, higher integrability assumptions on f are needed to give a meaning to the Vlasov–Poisson equation in the sense of distributions. For example, when $d = 3$, for the term Ef to belong to L_{loc}^1 one needs to have $f \in L^p$ with $p = (12 + 3\sqrt{5})/11$ (see for instance [34, 35]). To drop out this higher integrability hypotheses, in [34, 35] the authors considered the concept of renormalized solutions and obtained global existence provided that the total energy is finite and $f_0 \log(1 + f_0) \in L^1$. In addition, under some suitable integrability hypotheses on f , they can show that the concepts of weak and renormalized solutions are equivalent. For bounded density f , renormalization property holds because elliptic regularity of the Poisson equation leads to $E \in W^{1,p}$ with $p > 1$ (see [34, 35]). For the Vlasov–Maxwell system the only available global existence result is in [36], where the authors have constructed weak solutions for which it is not possible to show the renormalization property. Indeed, the best electromagnetic field regularity,

obtained so far for the DiPerna–Lions weak solutions, is in [22], where the authors show that the electromagnetic field (E, B) belongs to $H_{\text{loc}}^s(\mathbb{R}_*^+ \times \mathbb{R}^3)$, with $s = 6/(13 + \sqrt{142})$, if the macroscopic kinetic energy is in L^2 .

Regularity of the rough vector field considered above, i.e., Sobolev or BV vector fields, is somehow like the Lipschitz case because there is always a control (in Lebesgue spaces or in the space of measures with finite total variation) on an entire derivative of the vector field. By contrast, when b is not Lipschitz-like, the use of the chain rule is no longer justified, and many counterexamples to renormalization have been obtained in [2, 4–7, 29–31, 33, 65].

Here, we show that the renormalization property holds for an electromagnetic field with only a fractional derivative in some Lebesgue spaces, i.e., $E, B \in L_t^\infty W_x^{\beta,q}$ with $0 < \beta < 1$ and $1 \leq q \leq \infty$. To compensate for this loss of derivative for the electromagnetic field, the density f requires additional smoothness, typically fractional Sobolev differentiability in phase-space, i.e., $f \in L_t^1 W_{x,\xi}^{\alpha,p}$ with $0 < \alpha < 1$ and $1 \leq p \leq \infty$. We determine Onsager-type exponents [44] α and β , which ensure conservation of all entropies and guarantee that the renormalization property holds. As concerns the conservation of total energy, if the macroscopic kinetic energy is in L^2 , we then show that total energy is preserved. A comparable work has been done in [3] for the renormalization of an active scalar transport equation.

A similar situation occurs with systems of conservation laws of continuum physics, which are endowed with natural companion laws: the so-called entropy conditions (inequality versus equality) coming from the second law of thermodynamics. In [20, 52] the authors have determined the critical regularity of weak solutions to a general system of conservation laws to satisfy an associated entropy conservation law as an equality. They obtained the famous Onsager exponent $1/3$ [58]. The first result of this kind was obtained in [32] (see also [43]), where the authors have shown that weak solutions of the incompressible Euler equations conserve energy provided they possess fractional Besov differentiability of order greater than $1/3$. Such a result has been extended in various directions: In [27, 41, 47] the Besov criterium has been optimized; in [39, 45, 52, 56, 67] the authors have considered compressible Euler, Navier-Stokes, and magnetohydrodynamic equations; works [18, 19, 40, 60, 61] include boundary effects.

2. Basic properties. In this section, we recall the basic properties of the relativistic Vlasov–Maxwell system, which are valid for any smooth solution (f, E, B) , vanishing at infinity. These formal properties, in particular, natural a priori estimates, are the key cornerstones for proving the local-in-time well-posedness of this system [48]. Consider the following set of equations:

$$\partial_t f + v \cdot \nabla_x f + (E + v \times B) \cdot \nabla_\xi f = 0, \quad (t, x, \xi) \in \mathbb{R}^+ \times \mathbb{R}^3 \times \mathbb{R}^3, \quad (11)$$

$$\partial_t E - \nabla \times B = -j, \quad \partial_t B + \nabla \times E = 0, \quad (12)$$

$$\rho(t, x) = \int_{\mathbb{R}^3} f(t, x, \xi) d\xi, \quad j(t, x) = \int_{\mathbb{R}^3} v f(t, x, \xi) d\xi, \quad (13)$$

$$\gamma = \sqrt{1 + |\xi|^2}, \quad v = \nabla_\xi \gamma = \frac{\xi}{\sqrt{1 + |\xi|^2}}. \quad (14)$$

Observe that once the current density j and initial data (E_0, B_0) are given, the Maxwell equations (12) are well defined. Indeed the Maxwell operator M defined by,

$$X \mapsto MX = \begin{pmatrix} -\nabla \times B \\ \nabla \times E \end{pmatrix} \quad \text{with} \quad X = \begin{pmatrix} E \\ B \end{pmatrix}, \quad (15)$$

is the generator of a strongly continuous unitary group $t \mapsto S(t) := \exp(-Mt)$ in $L^2(\mathbb{R}^3)$ [14, 42, 66]. If $(E(t=0), B(t=0)) = (E_0, B_0) \in L^2(\mathbb{R}^3)$ and $j \in L^1(\mathbb{R}^+; L^2(\mathbb{R}^3))$, then, using the properties of the group $S(t)$ and the Duhamel formula, we can show that the solution (E, B) to (12) belongs to $\mathcal{C}(\mathbb{R}^+; L^2(\mathbb{R}^3))$. Moreover for any $s \geq 0$, the H^s regularity is preserved, i.e., the previous statement remains valid if we replace $L^2(\mathbb{R}^3)$ by $H^s(\mathbb{R}^3)$. In the same way, once the smooth electromagnetic field (E, B) and initial data $f_0(x, \xi)$ are given, the Vlasov equation is then well defined. Indeed, introducing the characteristic curves $t \mapsto (X(t), \Xi(t))$, which are the unique and smooth solution to the ODEs,

$$\frac{dX}{dt}(t) = v(\Xi(t)), \quad \frac{d\Xi}{dt}(t) = E(t, X(t)) + v(\Xi(t)) \times B(t, X(t)), \quad (16)$$

$$X(0; 0, x, \xi) = x, \quad \Xi(0; 0, x, \xi) = \xi, \quad (17)$$

the Lagrangian solution to (11) is given by (e.g., see [24])

$$f(t, x, \xi) = f_0(X(0; t, x, \xi), \Xi(0, t, x, \xi)). \quad (18)$$

The relativistic Vlasov–Maxwell system (11)-(14) satisfies some formal conservation laws, summarized in

PROPOSITION 1. Let (f, E, B) be a smooth solution, vanishing at infinity, to the relativistic Vlasov–Maxwell system (11)-(14). Then the following a priori estimates hold:

1. (Maximum principle). $0 \leq f_0 \leq M < \infty$ implies $0 \leq f(t) \leq M$ for all $t > 0$.
2. (L^p -norm conservation). For all $t \geq 0$, and $1 \leq p \leq \infty$, one has $\|f(t)\|_{L^p(\mathbb{R}^6)} = \|f_0\|_{L^p(\mathbb{R}^6)}$.
3. (Entropies). For any function $\mathcal{H} \in \mathcal{C}^1(\mathbb{R}^+; \mathbb{R}^+)$, one has for all $t \geq 0$,

$$\frac{d}{dt} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \mathcal{H}(f(t)) d\xi dx = 0.$$

4. (Energy conservation). For all $t \geq 0$ one has

$$\frac{d}{dt} \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} (\gamma(\xi) - 1) f(t) d\xi dx + \frac{1}{2} \int_{\mathbb{R}^3} (|E(t)|^2 + |B(t)|^2) dx \right) = 0.$$

5. (Momentum conservation). For all $t \geq 0$ one has

$$\frac{d}{dt} \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \xi f(t) d\xi dx + \int_{\mathbb{R}^3} (E(t) \times B(t)) dx \right) = 0.$$

Proof. The proof is standard and can be found, for instance, in [24]. \square

REMARK 1. Properties of Proposition 1 are key ingredients to obtain the global-in-time existence of weak solutions [36, 48, 59] and the local-in-time existence, uniqueness, and stability of classical solutions (e.g., see [48] and the references therein). Properties of Proposition 1 are also independent of other a priori invariances described below. Indeed

from the Maxwell–Faraday equation, $\partial_t B + \nabla \times E = 0$, we deduce that $\partial_t \nabla \cdot B = 0$, which leads to $\nabla \cdot B(t) = 0$ for all $t > 0$, if initially $\nabla \cdot B_0 = 0$. In a similar way, from the Maxwell–Ampère equation, $\partial_t E - \nabla \times B = -j$, we deduce that $\partial_t(\nabla \cdot E) + \nabla \cdot j = 0$. Using the charge conservation law (7) (obtained by integration of the Vlasov equation (11) with respect to ξ) we then obtain $\partial_t(\rho - \nabla \cdot E) = 0$, which leads to $\nabla \cdot E(t) = \rho(t)$ for all $t > 0$, if initially $\nabla \cdot E_0 = \rho_0$.

3. Renormalization property and entropies conservation.

3.1. Notation. We denote by \mathbb{R}^+ the nonnegative real numbers, by $\mathcal{D}(\mathbb{R}^d)$ the space of indefinitely differentiable with compact support, and by $\mathcal{D}'(\mathbb{R}^d)$ the space of distributions. We also denote by $\mathcal{S}(\mathbb{R}^d)$ the space of indefinitely differentiable and rapidly decreasing functions, and by $\mathcal{S}'(\mathbb{R}^d)$ the dual of $\mathcal{S}(\mathbb{R}^d)$, i.e., the space of tempered distributions. We use the notation $B_{p,q}^\alpha$ ($0 < \alpha < 1$, $1 \leq p \leq \infty$, $1 \leq q \leq \infty$) for Besov spaces, the definition of which can be found, e.g., in [1, 21, 63, 64]. The notation $W^{\alpha,p}$ ($0 < \alpha < 1$, $1 \leq p \leq \infty$) stands for the generalized Sobolev spaces of fractional order, whose precise definition can also be found, e.g., in [1, 21, 63, 64]. Let us simply recall first $W^{\alpha,p}(\mathbb{R}^d) = B_{p,p}^\alpha(\mathbb{R}^d)$ for α positive but not an integer and $1 \leq p \leq \infty$, and second the continuous embeddings: $B_{p,1}^\alpha(\mathbb{R}^d) \subset W^{\alpha,p}(\mathbb{R}^d) \subset B_{p,\infty}^\alpha(\mathbb{R}^d)$ with $1 \leq p \leq \infty$. We also define the functional space L_γ^1 such that

$$L_\gamma^1 = \left\{ f \geq 0 \text{ a.e. } \mid \|f\|_{L_\gamma^1(\mathbb{R}^6)} := \int_{\mathbb{R}^6} \gamma f \, dx d\xi < +\infty \right\}. \quad (19)$$

Moreover we define the function space \mathcal{E} such that

$$\mathcal{E} = \left\{ \mathcal{H} : \mathbb{R}^+ \mapsto \mathbb{R}^+; \mathcal{H} \text{ is nondecreasing, } \mathcal{H} \in \mathcal{C}^1(\mathbb{R}^+; \mathbb{R}^+), \lim_{\sigma \rightarrow +\infty} \frac{\mathcal{H}(\sigma)}{\sigma} = +\infty \right\}. \quad (20)$$

3.2. Main theorems. In this section we present our main results. For this, we need to recall the DiPerna–Lions theorem, which is the only existing result concerning the existence of global-in-time (weak) solutions to the Vlasov–Maxwell system in \mathbb{R}^6 .

THEOREM 1 (DiPerna–Lions [36]). Let $f_0 \in L_\gamma^1 \cap L^\infty(\mathbb{R}^6)$, and $E_0, B_0 \in L^2(\mathbb{R}^3)$, be initial conditions which satisfy the constraints,

$$\nabla \cdot B_0 = 0, \quad \nabla \cdot E_0 = \rho_0 = \int_{\mathbb{R}^3} f_0 \, d\xi \quad \text{in } \mathcal{D}'(\mathbb{R}^3).$$

Then, there exists a global-in-time weak solution of the relativistic Vlasov–Maxwell system, i.e., there are

$$f \in L^\infty(\mathbb{R}^+; L_\gamma^1 \cap L^\infty(\mathbb{R}^6)), \quad E, B \in L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^3)), \quad \rho, j \in L^\infty(\mathbb{R}^+; L^{4/3}(\mathbb{R}^3)), \quad (21)$$

such that (f, E, B) satisfy (11)–(12) in the sense of distributions, with ρ, j defined in terms of (13). Constraint equation (3) and the charge conservation law (7) are satisfied in the sense of distributions.

In addition, the mapping $t \mapsto f(t)$ (resp., $t \mapsto (E(t), B(t))$) is continuous with respect to the following topologies: the standard topology in the space of distributions $\mathcal{D}'(\mathbb{R}^6)$

(resp., $\mathcal{D}'(\mathbb{R}^3)$), the weak topology of $L^2(\mathbb{R}^6)$ (resp., $L^2(\mathbb{R}^3)$), and the strong topology of $H^{-s}(\Omega)$ for any $s > 0$ and any bounded subset Ω of \mathbb{R}^6 (resp., \mathbb{R}^3).

Futhermore, the total mass,

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} f(t) dx d\xi,$$

is independent of time, and one has,

$\|f(t)\|_{L^p(\mathbb{R}^6)} = \|f_0\|_{L^p(\mathbb{R}^6)}$ a.e. $t \geq 0$ for $1 \leq p \leq +\infty$, and $\mathcal{E}(t) \leq \mathcal{E}_0 < \infty$ a.e. $t \geq 0$, with the definition,

$$\mathcal{E}(t) := \int_{\mathbb{R}^6} \gamma f(t) d\xi dx + \frac{1}{2} \int_{\mathbb{R}^3} (|E(t)|^2 + |B(t)|^2) dx. \quad (22)$$

REMARK 2. 1. The conservation of mass and all the L^p norms was in fact proved in [59].
2. Using lower semi-continuity, weak solutions of Theorem 1 satisfy, for all $\mathcal{H} \in \mathscr{C}^1(\mathbb{R}^+; \mathbb{R}^+)$,

$$\int_{\mathbb{R}^6} \mathcal{H}(f(t)) d\xi dx \leq \int_{\mathbb{R}^6} \mathcal{H}(f_0) d\xi dx \quad \text{for } t \geq 0.$$

Now we intend to produce supplementary sufficient regularity conditions, which will imply the validity of supplementary conservation laws. As the first step this is the aim of Theorem 2 below: indeed, we first give sufficient regularity hypotheses which couple the regularity of the distribution function f with the regularity of the electromagnetic field (E, B) . In the second step, we use Theorem 2 and the results of [22] on the regularity of DiPerna–Lions weak solutions, to obtain Corollary 1 below, which involves only a sufficient regularity condition on the distribution function f . As concerns the renormalization property and entropies conservation, we have

THEOREM 2. Let (f, E, B) be a weak solution of the relativistic Vlasov–Maxwell system (11)–(14), given by Theorem 1. Assume that with

$$\alpha, \beta \in \mathbb{R}, \quad 0 < \alpha, \beta < 1, \quad \text{and} \quad \alpha\beta + \beta + 3\alpha - 1 > 0, \quad (23)$$

this weak solution satisfies for some $(p, q) \in \mathbb{N}_*^2$ with

$$\begin{aligned} \frac{1}{p} + \frac{1}{q} = \frac{1}{r} \leq 1 &\quad \text{if } 1 \leq p, q < \infty, \\ 1 \leq r < \infty &\quad \text{is arbitrary if } p = q = \infty, \end{aligned} \quad (24)$$

the supplementary regularity hypotheses,

$$f \in L^\infty(0, T; W^{\alpha, p}(\mathbb{R}^6)) \quad \text{and} \quad E, B \in L^\infty(0, T; W^{\beta, q}(\mathbb{R}^3)). \quad (25)$$

Then for any entropy function $\mathcal{H} \in \mathscr{C}^1(\mathbb{R}^+; \mathbb{R}^+)$, we have the renormalization property,

$$\partial_t(\mathcal{H}(f)) + \nabla_x \cdot (v\mathcal{H}(f)) + \nabla_\xi \cdot (F_L \mathcal{H}(f)) = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^6). \quad (26)$$

Moreover, if $\mathcal{H} \in \mathscr{C}$ and the map,

$$t \mapsto f(t, \cdot, \cdot) \text{ is uniformly integrable in } \mathbb{R}^6 \text{ for a.e. } t \in [0; T], \quad (27)$$

then we have the local entropy conservation laws,

$$\partial_t \left(\int_{\mathbb{R}^3} d\xi \mathcal{H}(f) \right) + \nabla_x \cdot \left(\int_{\mathbb{R}^3} d\xi v \mathcal{H}(f) \right) = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^3), \quad (28)$$

$$\partial_t \left(\int_{\mathbb{R}^3} dx \mathcal{H}(f) \right) + \nabla_\xi \cdot \left(\int_{\mathbb{R}^3} dx F_L \mathcal{H}(f) \right) = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^3), \quad (29)$$

and the global entropy conservation law,

$$\int_{\mathbb{R}^6} \mathcal{H}(f(t, x, \xi)) d\xi dx = \int_{\mathbb{R}^6} \mathcal{H}(f(s, x, \xi)) d\xi dx \quad \text{for } 0 < s \leq t < T. \quad (30)$$

The proof of Theorem 2 is postponed to Section 3.3. A few remarks are now in order.

REMARK 3. In fact, Theorem 2 is also true for the Vlasov–Poisson and the nonrelativistic Vlasov–Maxwell systems, under the same regularity assumptions.

REMARK 4. 1. In fact, Theorem 2 still holds when we replace Sobolev spaces $W^{\alpha,p}$ (resp., $W^{\beta,q}$) by Besov spaces $B_{p,\infty}^{\alpha+\epsilon}$ (resp., $B_{q,\infty}^\beta$) with $\epsilon > 0$. Indeed, even if Besov spaces $B_{p,\infty}^\alpha$ do not share the restriction property (needed for proving commutator estimates of Lemma 2), we still have the following result (see [15, 26, 54]): let $N \geq 2$, $1 \leq d < N$, $0 < p < q \leq \infty$, $\alpha' > N(1/p - 1)_+$, and $f \in B_{p,q}^{\alpha'}(\mathbb{R}^N)$. Then,

$$f(\cdot, y) \in \bigcap_{\alpha < \alpha'} B_{p,q}^\alpha(\mathbb{R}^d) \quad \text{for a.e. } y \in \mathbb{R}^{N-d}.$$

Therefore, in the Besov spaces framework, replacing α by $\alpha + \epsilon$ with $\epsilon > 0$ in (23), we observe that the condition $\alpha\beta + \beta + 3\alpha - 1 > 0$ keeps the same, whereas the phase-space regularity of f is slightly better than $B_{p,\infty}^\alpha$. Since the interpolation between $B_{p,\infty}^{\alpha+\epsilon}$ and $B_{p,p}^\alpha$ is $B_{p,r}^{\alpha'}$, with $\alpha < \alpha' < \alpha + \epsilon$, and $1 \leq r \leq \infty$, (e.g., Theorem 6.4.5 in [21]), we then have $B_{p,\infty}^{\alpha+\epsilon} \subset W^{\alpha,p}$.

2. Theorem 2 also includes the Hölder spaces where

$$f \in L^\infty(0, T; \mathscr{C}^{0,\alpha}(\mathbb{R}^6)) \quad \text{and} \quad E, B \in L^\infty(0, T; \mathscr{C}^{0,\beta}(\mathbb{R}^3)).$$

It corresponds to the case where $p = q = \infty$ (24)–(25) $\mathscr{C}^{0,\alpha} = W^{\alpha,\infty}$.

REMARK 5. Our result is almost in agreement with the structure-function scaling exponents derived in the study of dissipative anomalies in nearly collisionless plasma turbulence [44].

1. Here, the rigorous analysis is purely deterministic and regularity conditions (23)–(25) give a sufficient condition for the conservation of entropies for any individual solution as in [44]. In other words, by contraposition, a necessary condition for anomalous dissipation/nonconservation of entropies is, $\alpha\beta + \beta + 3\alpha - 1 < 0$ with $0 < \alpha, \beta < 1$. Nevertheless, this condition is not sufficient. Indeed, as in fluid mechanics with the Onsager critical regularity exponent $1/3$ [16, 17, 62], this necessary condition does not rule out the existence of some solutions that are less regular than the critical regularity (exponent) and that also satisfy the absence of anomalous entropy dissipation.

2. In [44] the author obtains, in a particular case, the critical exponent value $\alpha = \sqrt{5} - 2$, assuming that $f \in L^\infty(0, T; B_{p,\infty}^\alpha(\mathbb{R}_x^3; B_{p,\infty}^\alpha(\mathbb{R}_\xi^3)))$ and $E, B \in L^\infty(0, T; B_{p,\infty}^\alpha(\mathbb{R}^3))$ with $p \geq 3$. From Remark 4 on the restriction property of Besov spaces, in order to obtain $f \in L^\infty(0, T; B_{p,\infty}^\alpha(\mathbb{R}_x^3; B_{p,\infty}^\alpha(\mathbb{R}_\xi^3)))$, we must require the distribution function f to belong to the functional space $L^\infty(0, T; B_{p,\infty}^{\alpha+\epsilon}(\mathbb{R}^6))$ with $\epsilon > 0$. Now, taking $\alpha = \beta$, the condition $\alpha\beta + \beta + 3\alpha - 1 > 0$ in (23) becomes $\alpha^2 + 4\alpha - 1 > 0$, which is satisfied for $\alpha > \sqrt{5} - 2$. We then recover the same critical exponent value $\alpha = \sqrt{5} - 2$, but for $f \in L(0, T; W^{p,\alpha}(\mathbb{R}^6))$ and $E, B \in L^\infty(0, T; W^{q,\alpha}(\mathbb{R}^3))$ with $1/p + 1/q \leq 1$. Therefore, our regularity conditions (23)-(25) are weaker, but less restrictive than those of [44]. Indeed, we have $B_{p,\infty}^{\alpha+\epsilon} \subset W^{\alpha,p}$, $\forall \epsilon > 0$, and the condition $1/p + 1/q \leq 1$ is less restrictive than the condition $p = q \geq 3$.
3. In [44] the author obtains a refined version of the condition (23), by considering anisotropic regularity for the distribution function f between the space of velocities and the physical space, namely, $f \in L^\infty(0, T; B_{p,\infty}^\kappa(\mathbb{R}_x^3; B_{p,\infty}^\sigma(\mathbb{R}_\xi^3)))$. From Remark 4 on the restriction property of Besov spaces, this anisotropic regularity implies that $f \in L^\infty(0, T; B_{p,\infty}^{\alpha+\epsilon}(\mathbb{R}^6))$ with $\alpha := \max\{\kappa, \sigma\}$ and $\epsilon > 0$. This regularity condition is still more restrictive than our regularity condition, namely, $f \in L^\infty(0, T; W^{\alpha,p}(\mathbb{R}^6))$ with the same index α . In addition, anisotropic regularity in phase space is questionable because of the following physical argument. Phase-space turbulence involves typical structures known as vortices that are the result of the filamentation and the trapping (or wave-particle synchronization) phenomena. The fact that characteristic curves roll up in phase space seems to contradict that phase-space regularity is anisotropic between the space of velocities and the physical space. On the contrary, this mixing motion must propagate regularity versus singularities from one direction to another. By contrast, anisotropic regularity between the electromagnetic field (E, B) and the distribution f is justified and crucial, because the velocity integration of f can lead to additional regularity in the physical space for the moments such as charge and current densities, and hence for the electromagnetic field (through Maxwell's equations). This is the essence of averaging lemma [36] and the spirit of regularity results obtained for the DiPerna–Lions weak solutions [22, 25]. This anisotropy of regularity is handled both here and in [44].

REMARK 6. In the nonself-consistent case, i.e., when the Lorentz force F_L is a given external force, renormalization property (26) implies straightforwardly the uniqueness of weak solutions of Theorem 2, if such solutions exist. Indeed, let f^i , $i = 1, 2$, be two solutions of the Vlasov equation (11), with initial conditions f_0^i , $i = 1, 2$, and where the electromagnetic field (E, B) is prescribed. Such solutions satisfy the regularity properties of Theorem 2, in particular (25). Setting $f = f^1 - f^2$, and taking $\mathcal{H}(\cdot) = (\cdot)^2$ ($\mathcal{H} \in \mathcal{E}$), we obtain from Theorem 2,

$$\partial_t(\mathcal{H}(f)) + \nabla_x \cdot (v\mathcal{H}(f)) + \nabla_\xi \cdot (F_L \mathcal{H}(f)) = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^6),$$

and

$$\int_{\mathbb{R}^6} \mathcal{H}(f(t)) d\xi dx = \int_{\mathbb{R}^6} \mathcal{H}(f_0) d\xi dx.$$

Therefore, taking $f_0^1 = f_0^2$, i.e., $f_0 = f^1 - f^2 = 0$, we obtain $f = 0$ a.e., i.e., $f_1 = f_2$ a.e. In a similar way we can show the following comparison principle: $f_0^1 \leq f_0^2$ a.e. implies $f^1 \leq f^2$ a.e. Two open issues remain. The first one is the uniqueness of solutions of Theorem 2, which corresponds to the self-consistent case. Of course the existence of solutions of Theorem 2 is also an open big problem. Following the program of [36], the second one is the existence and uniqueness of corresponding Lagrangian solutions, i.e., solutions constructed from almost-everywhere-well-defined characteristic curves, as in the smooth framework (16)-(18).

REMARK 7. Another open issue is the case of bounded domains in space, with specular reflection and/or absorbing conditions [49]. This is not an easy task since, for such natural boundary conditions, some singularities could occur at the boundary and propagate inside the domain [50, 51].

From Theorem 2 and the result of [22] on the regularity of the DiPerna–Lions weak solutions, we deduce the following corollary, which involves hypotheses concerning only the distribution function f .

COROLLARY 1. Let $\beta = 6/(13 + \sqrt{142})$, and let $\alpha \in \mathbb{R}$ be the solution to

$$\alpha\beta + \beta + 3\alpha - 1 > 0 \quad \text{and} \quad 0 < \alpha < 1. \quad (31)$$

Let (f, E, B) be a weak solution to the relativistic Vlasov–Maxwell system (11)–(14), given by Theorem 1. Assume the additional hypotheses: initial conditions (E_0, B_0) belong to $H^1(\mathbb{R}^3)$, the distribution function f satisfies the supplementary integrability condition,

$$\int_{\mathbb{R}^3} \gamma f d\xi \in L^\infty(0, T; L^2(\mathbb{R}^3)), \quad (32)$$

and the regularity assumption,

$$f \in L^\infty(0, T; H^\alpha(\mathbb{R}^6)). \quad (33)$$

Then for any entropy function $\mathcal{H} \in \mathscr{C}^1(\mathbb{R}^+; \mathbb{R}^+)$, renormalization property (26) holds. Moreover, if $\mathcal{H} \in \mathcal{E}$ and the map $t \mapsto f(t, \cdot, \cdot)$ is uniformly integrable in \mathbb{R}^6 for a.e. $t \in \mathbb{R}^+$, then local entropy conservation laws (28)–(29), as well as global entropy conservation law (30), hold.

Proof of Corollary 1. Using assumptions $E_0, B_0 \in H^1(\mathbb{R}^3)$, and (32) from Theorem 1.1 of [22], we obtain that the electromagnetic field (E, B) belongs to $H_{\text{loc}}^\beta(\mathbb{R}^+ \times \mathbb{R}^3)$ with $\beta = 6/(13 + \sqrt{142})$. Setting $p = q = 2$ and $\beta = 6/(13 + \sqrt{142})$ in the hypotheses of Theorem 2, and using assumption (33) under constraints (31), we obtain from Theorem 2 the desired result. \square

REMARK 8. From Corollary 1, we deduce that for α such that

$$1 > \alpha > \frac{1 - \beta}{3 + \beta} = \frac{7 + \sqrt{142}}{45 + 3\sqrt{142}} \simeq 0.234, \quad (34)$$

the Vlasov equation (11), which is a first-order conservation law in the phase-space \mathbb{R}^6 , has an infinite number of conserved entropies. A similar situation occurs with general systems of conservation laws, which are studied in [20] within the regularity framework of Hölder spaces $\mathcal{C}^{0,\alpha}$. Nevertheless, in [20], the authors show conservation of entropies under the sufficient condition $\alpha > 1/3$ (the famous Onsager exponent [58]), which is more restrictive than the present result, from two points of view. First, Sobolev spaces H^α are less regular than Hölder spaces $\mathcal{C}^{0,\alpha}$ for the same α . Secondly, our index α is smaller than $1/3$. An explanation of such a discrepancy comes from our commutator estimates which exploit the anisotropy between the velocity and physical spaces, whereas commutator estimates in [20, 52] use some Taylor expansions, which does not advantage a particular direction of space. Finally, we observe that the critical exponent $\alpha = (7 + \sqrt{142})/(45 + 3\sqrt{142})$, which is smaller than $\sqrt{5} - 2$, cannot be retrieved with the method of [44], since the latter is obtained under the condition $p = q \geq 3$ and hence cannot deal with the case $p = q = 2$.

3.3. Proof of Theorem 2. Before giving the proof of Theorem 2, we first introduce some standard regularization operators and we recall their main properties. Using a smooth nonnegative function ϱ such that

$$\tau \mapsto \varrho(\tau) \geq 0, \quad \varrho \in \mathcal{D}(\mathbb{R}), \quad \text{supp}(\varrho) \subset]-1, 1[, \quad \int_{\mathbb{R}} \varrho(\tau) d\tau = 1, \quad (35)$$

one defines the radially-symmetric compactly-supported Friedrichs mollifier $z \mapsto \varrho_\epsilon(z)$, given by

$$\begin{aligned} \mathbb{R}^d &\longrightarrow \mathbb{R}^+ \\ z &\longmapsto \varrho_\epsilon(z) = \frac{1}{\epsilon^d} \varrho\left(\frac{|z|}{\epsilon}\right), \quad \epsilon > 0. \end{aligned} \quad (36)$$

For any distribution $f \in \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^6)$, we define its \mathcal{C}^∞ -regularization by

$$f^{\eta, \varepsilon, \delta}(t, x, \xi) = \varrho_\eta(t) *_t \varrho_\varepsilon(x) *_x \varrho_\delta(\xi) *_\xi f(t, x, \xi), \quad (37)$$

where the operator $*$ denotes the standard convolution product. We denote by $\langle \cdot, \cdot \rangle$ the dual bracket between spaces \mathcal{D}' and \mathcal{D} . Using previous definitions, we have for the regularization operator $(\cdot)^\epsilon$ the following standard properties (see, e.g., [8]), which are summarized in the following.

LEMMA 1. 1. For any distribution $f \in \mathcal{D}'(\mathbb{R}^d)$, we have

$$\langle f^\epsilon, g \rangle = \langle f, g^\epsilon \rangle, \quad g \in \mathcal{D}(\mathbb{R}^d).$$

2. For any function $f \in L^1 \cap L^\infty \cap W^{\alpha,p}(\mathbb{R}^d)$, with $0 < \alpha < 1$ and $1 \leq p \leq \infty$, there exists a constant C such that

$$\begin{aligned}\|f^\epsilon\|_{L^q(\mathbb{R}^d)} &\leq \|f\|_{L^q(\mathbb{R}^d)}, \quad 1 \leq q \leq \infty, \\ \|f^\epsilon\|_{W^{\alpha,p}(\mathbb{R}^d)} &\leq \|f\|_{W^{\alpha,p}(\mathbb{R}^d)}, \\ \|f^\epsilon - f\|_{L^p(\mathbb{R}^d)} &\leq C\epsilon^\alpha \|f\|_{W^{\alpha,p}(\mathbb{R}^d)}, \\ \|\nabla f^\epsilon\|_{L^p(\mathbb{R}^d)} &\leq C\epsilon^{\alpha-1} \|f\|_{W^{\alpha,p}(\mathbb{R}^d)}.\end{aligned}$$

3. For any function $f \in B_{p,\infty}^\alpha(\mathbb{R}^d)$, with $0 < \alpha < 1$ and $1 \leq p \leq \infty$, there exists a constant C such that

$$\|f(\cdot - z) - f(\cdot)\|_{L^p(\mathbb{R}^d)} \leq C|z|^\alpha \|f\|_{B_{p,\infty}^\alpha(\mathbb{R}^d)}.$$

Proof. Since the proof is elementary, it is left to the reader. \square

In order to prove Theorem 2, we use some commutator estimates which are given by

LEMMA 2. Let (f, E, B) be a weak solution of the relativistic Vlasov–Maxwell system (11)–(14), given by Theorem 1, satisfying the regularity assumptions (23)–(25) of Theorem 2. Let us recall that $F_L := E + v \times B$ is the Lorentz force field. Then there exist a constant C_{fs} depending on $\|f\|_{L^\infty(0,T;W^{\alpha,p}(\mathbb{R}^6))}$ and a constant C_{fl} depending on $\|f\|_{L^\infty(0,T;W^{\alpha,p}(\mathbb{R}^6))}$, $\|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))}$, and $\|B\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))}$ such that

$$\|\nabla_x \cdot ((vf)^{\eta,\varepsilon,\delta} - v^\delta f^{\eta,\varepsilon,\delta})\|_{L^1(0,T;L^p(\mathbb{R}^6))} \leq C_{fs} \delta^{\alpha+1} \varepsilon^{\alpha-1} \quad (38)$$

and

$$\|\nabla_\xi \cdot ((F_L f)^{\eta,\varepsilon,\delta} - F_L^{\eta,\varepsilon,\delta} f^{\eta,\varepsilon,\delta})\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \leq C_{fl} (\varepsilon^{\beta+\alpha} \delta^{\alpha-1} + \delta^\alpha), \quad (39)$$

where α, β, p, q , and r satisfy relations (23)–(24).

REMARK 9. The precise estimates obtained in Lemma 2 seem to be compulsory to obtain the precise Onsager exponents α, β in the main theorem, Theorem 2, instead of the general exponent 1/3 established for fluid models.

Proof. We start with two basic estimates, which will be used often throughout the proof. Using the fundamental theorem of calculus and

$$|\nabla_\xi v| = \left| \frac{I_3}{(1 + |\xi|^2)^{1/2}} - \frac{\xi \otimes \xi}{(1 + |\xi|^2)^{3/2}} \right| \leq \frac{2}{\sqrt{1 + |\xi|^2}} \leq 2, \quad (40)$$

we obtain the first basic estimate,

$$|v(\xi - w) - v(\xi)| \leq |w| \int_0^1 |\nabla v(\xi - \tau w)| d\tau \leq 2|w|. \quad (41)$$

Using the fundamental theorem of calculus twice, we obtain componentwise,

$$\begin{aligned}
v_i - v_i^\delta &= \int_{\mathbb{R}^3} dw \varrho_\delta(w) (v_i(\xi) - v_i(\xi - w)) \\
&= \sum_j \int_{\mathbb{R}^3} dw \varrho_\delta(w) w_j \int_0^1 d\tau \partial_j v_i(\xi - \tau w) \\
&= \sum_j \partial_j v_i(\xi) \int_{\mathbb{R}^3} dw \varrho_\delta(w) w_j \\
&\quad + \sum_{j,k} \int_{\mathbb{R}^3} dw \varrho_\delta(w) w_j w_k \int_0^1 d\tau \int_0^1 ds \partial_{jk}^2 v_i(\xi - s\tau w). \tag{42}
\end{aligned}$$

Since the smooth function ϱ is radially symmetric and compactly supported, we have,

$$\int_{\mathbb{R}^3} dw \varrho(w) w_i = 0 \quad \text{and} \quad \int_{\mathbb{R}^3} dw \varrho(w) |w_i| |w_j| \leq C_\varrho < +\infty \quad \forall i, j \in \{1, 2, 3\}, \tag{43}$$

where C_ϱ is a numerical constant depending only on the function ϱ . Using the first equality in (43), the first term of the right-hand side of (42) vanishes. Using the second inequality of (43), and

$$|\nabla_{jk}^2 v_i(\xi)| = \left| \frac{\delta_{ij}\xi_k}{(1+|\xi|^2)^{3/2}} + \frac{\delta_{jk}\xi_i}{(1+|\xi|^2)^{3/2}} + \frac{\delta_{ik}\xi_j}{(1+|\xi|^2)^{3/2}} - \frac{3\xi_i\xi_j\xi_k}{(1+|\xi|^2)^{3/2}} \right| \leq \frac{6}{1+|\xi|^2} \leq 6,$$

we obtain from (42) the second basic estimate,

$$|v - v^\delta| \leq 6C_\varrho \delta^2. \tag{44}$$

We now deal with commutator estimate (38) for the free-streaming term. We define

$$\begin{aligned}
r_{\eta,\varepsilon,\delta}(f, g)(t, x, \xi) &= \int_{\mathbb{R}} d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) \varrho_\delta(w) \\
&\quad (f(t - \tau, x - y, \xi - w) - f(t, x, \xi))(g(t - \tau, x - y, \xi - w) - g(t, x, \xi)). \tag{45}
\end{aligned}$$

Using (45), it is easy to check that

$$(vf)^{\eta,\varepsilon,\delta} = v^\delta f^{\eta,\varepsilon,\delta} + r_{\eta,\varepsilon,\delta}(v, f) - (f - f^{\eta,\varepsilon,\delta})(v - v^\delta). \tag{46}$$

Observing that

$$r_{\eta,\varepsilon,\delta}(v, f) = r_\delta(v, f^{\eta,\varepsilon}) + (f - f^{\eta,\varepsilon})(v - v^\delta),$$

equation (46) becomes

$$(vf)^{\eta,\varepsilon,\delta} - v^\delta f^{\eta,\varepsilon,\delta} = r_\delta(v, f^{\eta,\varepsilon}) - ((f^{\eta,\varepsilon})^\delta - f^{\eta,\varepsilon})(v - v^\delta). \tag{47}$$

Using estimate (41), Lemma 1, continuous embedding $W^{\alpha,p}(\mathbb{R}^d) \subset B_{p,\infty}^\alpha(\mathbb{R}^d)$ with $1 \leq p \leq \infty$, the restriction property for Sobolev spaces $W^{\alpha,p}(\mathbb{R}^d)$ (see Remark 4),

and regularity assumptions (23)-(25), we obtain,

$$\begin{aligned}
& \|\nabla_x \cdot r_\delta(v, f^{\eta, \varepsilon})\|_{L^1(0, T; L^p(\mathbb{R}^6))} \\
& \leq \int_0^T dt \int_{\mathbb{R}^3} dw \varrho_\delta(w) \|(v(\xi - w) - v(\xi)) \\
& \quad \cdot (\nabla_x f^{\eta, \varepsilon}(t, x, \xi - w) - \nabla_x f^{\eta, \varepsilon}(t, x, \xi))\|_{L^p(\mathbb{R}_{x\xi}^6)} \\
& \leq C \int_0^T dt \int_{\mathbb{R}^3} dw \varrho_\delta(w) |w|^{\alpha+1} \|\nabla_x f^{\eta, \varepsilon}(t)\|_{L^p(\mathbb{R}_x^3; B_{p, \infty}^\alpha(\mathbb{R}_\xi^3))} \\
& \leq C \delta^{\alpha+1} \int_0^T dt \|\nabla_x f^{\eta, \varepsilon}(t)\|_{L^p(\mathbb{R}_x^3; W^{\alpha, p}(\mathbb{R}_\xi^3))} \\
& \leq C \varepsilon^{\alpha-1} \delta^{\alpha+1} \int_0^T dt \|f^\eta(t)\|_{W^{\alpha, p}(\mathbb{R}_x^3; W^{\alpha, p}(\mathbb{R}_\xi^3))} \\
& \leq C \varepsilon^{\alpha-1} \delta^{\alpha+1} \int_0^T dt \varrho_\eta(t) * \|f(t)\|_{W^{\alpha, p}(\mathbb{R}^6)} \\
& \leq C \varepsilon^{\alpha-1} \delta^{\alpha+1} \|f\|_{L^1(0, T; W^{\alpha, p}(\mathbb{R}^6))}.
\end{aligned} \tag{48}$$

Using estimate (44), Lemma 1, the restriction property for Sobolev spaces $W^{\alpha, p}(\mathbb{R}^d)$, and regularity assumptions (23)-(25), we obtain,

$$\begin{aligned}
& \|\nabla_x \cdot ((f^{\eta, \varepsilon})^\delta - f^{\eta, \varepsilon})(v - v^\delta)\|_{L^1(0, T; L^p(\mathbb{R}^6))} \\
& \leq \|v - v^\delta\|_{L^\infty(\mathbb{R}^3)} \|(\nabla_x f^{\eta, \varepsilon})^\delta - \nabla_x f^{\eta, \varepsilon}\|_{L^1(0, T; L^p(\mathbb{R}^6))} \\
& \leq C \delta^{\alpha+1} \|\nabla_x f^{\eta, \varepsilon}\|_{L^1(0, T; L^p(\mathbb{R}_x^3; W^{\alpha, p}(\mathbb{R}_\xi^3)))} \\
& \leq C \varepsilon^{\alpha-1} \delta^{\alpha+1} \|f^\eta\|_{L^1(0, T; W^{\alpha, p}(\mathbb{R}_x^3; W^{\alpha, p}(\mathbb{R}_\xi^3)))} \\
& \leq C \varepsilon^{\alpha-1} \delta^{\alpha+1} \|f\|_{L^1(0, T; W^{\alpha, p}(\mathbb{R}^6))}.
\end{aligned} \tag{49}$$

Using (48)-(49), we obtain from (47), commutator estimate (38). We continue with commutator estimate (39) for the Lorentz force term. Using definition (45), we first make the following decomposition:

$$(F_L f)^{\eta, \varepsilon, \delta} - F_L^{\eta, \varepsilon, \delta} f^{\eta, \varepsilon, \delta} = T_E + T_B, \tag{50}$$

where

$$T_E = (E f^\delta)^{\eta, \varepsilon} - E^{\eta, \varepsilon} (f^\delta)^{\eta, \varepsilon} = r_{\eta, \varepsilon}(E, f^\delta) - (E - E^{\eta, \varepsilon})(f^\delta - (f^\delta)^{\eta, \varepsilon}) \tag{51}$$

and

$$T_B = (v \times B f)^{\varepsilon, \delta} - v^\delta \times B^\varepsilon f^{\varepsilon, \delta}. \tag{52}$$

Let us first deal with the term T_E . Passing to the limit $\eta \rightarrow 0$ in $r_{\eta, \varepsilon}(E, f^\delta)$, which can be justified by the Lebesgue dominated convergence theorem and regularity assumptions

(23)-(25), we obtain,

$$\begin{aligned} \|\nabla_\xi \cdot r_{\eta,\varepsilon}(E, f^\delta)\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} &\leq \|\nabla_\xi \cdot r_\varepsilon(E, f^\delta)\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \\ &\leq \int_{\mathbb{R}^3} dy \varrho_\varepsilon(y) \|E(t, x-y) - E(t, x)\| \\ &\quad \cdot (\nabla_\xi f^\delta(t, x-y, \xi) - \nabla_\xi f^\delta(t, x, \xi)) \|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))}. \end{aligned} \quad (53)$$

Using Hölder inequality, Lemma 1, continuous embedding $W^{\alpha,p}(\mathbb{R}^d) \subset B_{p,\infty}^\alpha(\mathbb{R}^d)$ with $1 \leq p \leq \infty$, the restriction property for Sobolev spaces $W^{\alpha,p}(\mathbb{R}^d)$, and regularity assumptions (23)-(25), we obtain from (53),

$$\begin{aligned} &\|\nabla_\xi \cdot r_{\eta,\varepsilon}(E, f^\delta)\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \\ &\leq \int_{\mathbb{R}^3} dy \varrho_\varepsilon(y) \|E(t, x-y) - E(t, x)\|_{L^\infty(0,T;L^q(\mathbb{R}_x^3))} \\ &\quad \|\nabla_\xi f^\delta(t, x-y, \xi) - \nabla_\xi f^\delta(t, x, \xi)\|_{L^1(0,T;L^p(\mathbb{R}_{x\xi}^6))} \\ &\leq C \int_{\mathbb{R}^3} dy \varrho_\varepsilon(y) |y|^{\alpha+\beta} \\ &\quad \|E\|_{L^\infty(0,T;B_{q,\infty}^\beta(\mathbb{R}^3))} \|\nabla_\xi f^\delta\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;B_{p,\infty}^\alpha(\mathbb{R}_x^3)))} \\ &\leq C\varepsilon^{\alpha+\beta} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|\nabla_\xi f^\delta\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;W^{\alpha,p}(\mathbb{R}_x^3)))} \\ &\leq C\varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}_\xi^3;W^{\alpha,p}(\mathbb{R}_x^3)))} \\ &\leq C\varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}^6))}. \end{aligned} \quad (54)$$

Using the Lebesgue dominated convergence theorem and regularity assumptions (23)-(25), we can pass to the limit $\eta \rightarrow 0$ in the term, $(E - E^{\eta,\varepsilon})(f^\delta - (f^\delta)^{\eta,\varepsilon})$, to obtain

$$\begin{aligned} &\|\nabla_\xi \cdot ((E - E^{\eta,\varepsilon})(f^\delta - (f^\delta)^{\eta,\varepsilon}))\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \\ &\leq \|\nabla_\xi \cdot ((E - E^\varepsilon)(f^\delta - (f^\delta)^\varepsilon))\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))}. \end{aligned} \quad (55)$$

Using Hölder inequality, Lemma 1, continuous embedding $W^{\alpha,p}(\mathbb{R}^d) \subset B_{p,\infty}^\alpha(\mathbb{R}^d)$ with $1 \leq p \leq \infty$, the restriction property for Sobolev spaces $W^{\alpha,p}(\mathbb{R}^d)$, and regularity assumptions (23)-(25), we obtain from (55),

$$\begin{aligned} &\|\nabla_\xi \cdot ((E - E^{\eta,\varepsilon})(f^\delta - (f^\delta)^{\eta,\varepsilon}))\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \\ &\leq \|E - E^\varepsilon\|_{L^\infty(0,T;L^q(\mathbb{R}^3))} \|\nabla_\xi f^\delta - (\nabla_\xi f^\delta)^\varepsilon\|_{L^1(0,T;L^p(\mathbb{R}^6))} \\ &\leq C\varepsilon^{\alpha+\beta} \|E\|_{L^\infty(0,T;B_{q,\infty}^\beta(\mathbb{R}^3))} \|\nabla_\xi f^\delta\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;B_{p,\infty}^\alpha(\mathbb{R}_x^3)))} \\ &\leq C\varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}_\xi^3;W^{\alpha,p}(\mathbb{R}_x^3)))} \\ &\leq C\varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}^6))}. \end{aligned} \quad (56)$$

From (53) and (56), we obtain,

$$\|\nabla_\xi \cdot T_E\|_{L^1(0,T;L^p(\mathbb{R}_x^3;L^r(\mathbb{R}_x^3)))} \leq C\varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|E\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}^6))}. \quad (57)$$

We now deal with the term T_B , given by (52), and which can be recast as,

$$\begin{aligned} T_B &= \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) \varrho_\delta(w) [v(\xi - w) - v(\xi)] \\ &\quad \times B(t - \tau, x - y) f(t - \tau, x - y, \xi - w) \\ &\quad + v \times [(Bf^\delta)^{\eta, \varepsilon} - B^{\eta, \varepsilon}(f^\delta)^{\eta, \varepsilon}] + (v - v^\delta) \times B^{\eta, \varepsilon}(f^\delta)^{\eta, \varepsilon} \\ &= T_{B1} + T_{B2} + T_{B3}. \end{aligned} \quad (58)$$

The term $\nabla_\xi \cdot T_{B1}$ can be decomposed as

$$\begin{aligned} \nabla_\xi \cdot T_{B1} &= \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) \nabla_w \varrho_\delta(w) \\ &\quad \cdot ([v(\xi - w) - v(\xi)] \times B(t - \tau, x - y)) f(t - \tau, x - y, \xi) \\ &\quad + \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) \nabla_w \varrho_\delta(w) \\ &\quad \cdot ([v(\xi - w) - v(\xi)] \times B(t - \tau, x - y)) \\ &\quad (f(t - \tau, x - y, \xi - w) - f(t - \tau, x - y, \xi)) \\ &= T_{B11} + T_{B12}. \end{aligned} \quad (59)$$

Using integration by parts, we observe that

$$\begin{aligned} T_{B11} &= \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) \varrho_\delta(w) \\ &\quad \nabla_w \cdot ([v(\xi - w) - v(\xi)] \times B(t - \tau, x - y)) f(t - \tau, x - y, \xi) = 0, \end{aligned} \quad (60)$$

because $\nabla_w \cdot ([v(\xi - w) - v(\xi)] \times B(t - \tau, x - y)) = 0$. Using Hölder inequality, estimate (41), Lemma 1, continuous embedding $W^{\alpha, p}(\mathbb{R}^d) \subset B_{p, \infty}^\alpha(\mathbb{R}^d)$ with $1 \leq p \leq \infty$, the restriction property for Sobolev spaces $W^{\alpha, p}(\mathbb{R}^d)$, and regularity assumptions (23)-(25), we obtain

$$\begin{aligned} \|T_{B12}\|_{L^1(0, T; L^p(\mathbb{R}_\xi^3; L^r(\mathbb{R}_x^3)))} &\leq 2 \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) |\nabla_w \varrho_\delta(w)| |w| \\ &\quad \|B(t - \tau, x - y)(f(t - \tau, x - y, \xi - w) - f(t - \tau, x - y, \xi))\|_{L^1(0, T; L^p(\mathbb{R}_\xi^3; L^r(\mathbb{R}_x^3)))} \\ &\leq 2 \int_0^T d\tau \int_{\mathbb{R}^3} dy \int_{\mathbb{R}^3} dw \varrho_\eta(\tau) \varrho_\varepsilon(y) |\nabla_w \varrho_\delta(w)| |w| \\ &\quad \|B(t - \tau, x - y)\|_{L^\infty(0, T; L^q(\mathbb{R}_x^3))} \|f(t - \tau, x - y, \xi - w) - f(t - \tau, x - y, \xi)\|_{L^1(0, T; L^p(\mathbb{R}_{x\xi}^6))} \\ &\leq C \int_{\mathbb{R}^3} dw |\nabla_w \varrho_\delta(w)| |w|^{\alpha+1} \|B\|_{L^\infty(0, T; L^q(\mathbb{R}_x^3))} \|f\|_{L^1(0, T; L^p(\mathbb{R}_x^3; B_{p, \infty}^\alpha(\mathbb{R}_\xi^3)))} \\ &\leq C \delta^\alpha \|B\|_{L^\infty(0, T; W^{\beta, q}(\mathbb{R}^3))} \|f\|_{L^1(0, T; W^{\alpha, p}(\mathbb{R}^6))}. \end{aligned} \quad (61)$$

In a way similar to how we have obtained estimate (57) for $\nabla_\xi \cdot T_E$, we also obtain for $\nabla_\xi \cdot T_{B2}$,

$$\|\nabla_\xi \cdot T_{B2}\|_{L^1(0, T; L^p(\mathbb{R}_\xi^3; L^r(\mathbb{R}_x^3)))} \leq C \varepsilon^{\alpha+\beta} \delta^{\alpha-1} \|B\|_{L^\infty(0, T; W^{\beta, q}(\mathbb{R}^3))} \|f\|_{L^1(0, T; W^{\alpha, p}(\mathbb{R}^6))}. \quad (62)$$

Using estimate (44), Hölder inequality, and Lemma 1, we obtain for $\nabla_\xi \cdot T_{B3}$,

$$\begin{aligned} \|\nabla_\xi \cdot T_{B3}\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} &\leq \|(v - v^\delta) \times B^{\eta,\varepsilon} (\nabla_\xi f^\delta)^{\eta,\varepsilon}\|_{L^1(0,T;L^p(\mathbb{R}_\xi^3;L^r(\mathbb{R}_x^3)))} \\ &\leq C\delta^2 \|B^{\eta,\varepsilon}\|_{L^\infty(0,T;L^q(\mathbb{R}_x^3))} \|\nabla_\xi f^{\eta,\delta,\varepsilon}\|_{L^1(0,T;L^p(\mathbb{R}^6))} \\ &\leq C\delta^{\alpha+1} \|B\|_{L^\infty(0,T;L^q(\mathbb{R}_x^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}^6))}. \end{aligned} \quad (63)$$

Gathering estimates (60)-(63), we obtain from decompositions (58)-(59),

$$\|\nabla_\xi \cdot T_B\|_{L^1(0,T;L^p(\mathbb{R}_x^3;L^r(\mathbb{R}_x^3)))} \leq C(\varepsilon^{\alpha+\beta} \delta^{\alpha-1} + \delta^\alpha) \|B\|_{L^\infty(0,T;W^{\beta,q}(\mathbb{R}^3))} \|f\|_{L^1(0,T;W^{\alpha,p}(\mathbb{R}^6))}. \quad (64)$$

Eventually, from (57) and (64), we obtain commutator estimate (39), which ends the proof of Lemma 2 \square

Proof of Theorem 2. Let us now give the proof of the main theorem. The weak formulation for the Vlasov equation reads,

$$\int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi f (\partial_t \Psi + v \cdot \nabla_x \Psi + F_L \cdot \nabla_\xi \Psi) = 0, \quad \forall \Psi \in \mathcal{D}((0,T) \times \mathbb{R}^6) \quad (65)$$

with $F_L := E + v \times B$. Let us note that all integrals in (65) have a sense since for DiPerna–Lions weak solutions [36] we have $f \in L^\infty(0,T;L^2(\mathbb{R}^6))$, and $E, B \in L^\infty(0,T;L^2(\mathbb{R}^3))$. We choose in (65) the test function,

$$\Psi = \Psi_{\varepsilon,\delta} = (\mathcal{H}'(f^{\eta,\varepsilon,\delta})\Phi)^{\eta,\varepsilon,\delta} \in \mathcal{D}((0,T) \times \mathbb{R}^6) \quad (66)$$

with $\Phi \in \mathcal{D}((0,T) \times \mathbb{R}^6)$ and $\mathcal{H} \in \mathscr{C}^1(\mathbb{R}^+; \mathbb{R}^+)$. Using the first property of Lemma 1 and successive integrations by parts, we obtain from (65)-(66),

$$\begin{aligned} &\int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \left\{ \mathcal{H}(f^{\eta,\varepsilon,\delta})(\partial_t \Phi + v^\delta \cdot \nabla_x \Phi + F_L^{\eta,\varepsilon,\delta} \cdot \nabla_\xi \Phi) \right. \\ &\left. + \Phi \mathcal{H}'(f^{\eta,\varepsilon,\delta}) \left[\nabla_x \cdot ((vf)^{\eta,\varepsilon,\delta} - v^\delta f^{\eta,\varepsilon,\delta}) + \nabla_\xi \cdot ((F_L f)^{\eta,\varepsilon,\delta} - F_L^{\eta,\varepsilon,\delta} f^{\eta,\varepsilon,\delta}) \right] \right\} = 0 \end{aligned} \quad (67)$$

for all $\Phi \in \mathcal{D}((0,T) \times \mathbb{R}^6)$. We now establish the renormalized Vlasov equation (26). Using regularity assumptions (23)-(25), (66), and Lemmas 1 and 2, we obtain from (67),

$$\begin{aligned} &\left| \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta,\varepsilon,\delta})(\partial_t \Phi + v^\delta \cdot \nabla_x \Phi + F_L^{\eta,\varepsilon,\delta} \cdot \nabla_\xi \Phi) \right| \\ &\leq C_* (\delta^{\alpha+1} \varepsilon^{\alpha-1} + \varepsilon^{\alpha+\beta} \delta^{\alpha-1} + \delta^\alpha), \end{aligned} \quad (68)$$

where C_* depends on $\|f\|_{L^\infty(0,T;L^\infty(\mathbb{R}^6))}$, C_{fs} , C_{fl} , \mathcal{H} , and Φ . Balancing contributions coming from the free-streaming and Lorentz force terms in the right-hand side of (68), we obtain

$$\varepsilon^{\alpha-1} \delta^2 - \delta - \varepsilon^{\alpha+\beta} = 0, \quad (69)$$

and estimate (68) becomes

$$\left| \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta,\varepsilon,\delta})(\partial_t \Phi + v^\delta \cdot \nabla_x \Phi + F_L^{\eta,\varepsilon,\delta} \cdot \nabla_\xi \Phi) \right| \leq C_* \eta \quad (70)$$

with the definition

$$\eta := \varepsilon^{\alpha-1} \delta^{\alpha+1}.$$

Solving quadratic equation (69) in δ , the only positive solution is given by

$$\delta = \frac{1 + \sqrt{1 + 4\varepsilon^{2\alpha+\beta-1}}}{2\varepsilon^{\alpha-1}}.$$

Two cases are to be considered according to the value of α and β :

- i) $2\alpha + \beta - 1 < 0$. We then have $\delta \simeq \varepsilon^{(\beta+1)/2}$, and $\eta \simeq \varepsilon^{(\alpha\beta+\beta+3\alpha-1)/2} \rightarrow 0$ as $\varepsilon \rightarrow 0$ if $\alpha\beta + \beta + 3\alpha - 1 > 0$.
- ii) $2\alpha + \beta - 1 \geq 0$. We then have $\delta \simeq \varepsilon^{1-\alpha}$, and $\eta \simeq \varepsilon^{\alpha(1-\alpha)} \rightarrow 0$ as $\varepsilon \rightarrow 0$ if $0 < \alpha < 1$.

Assuming that the free-streaming contribution dominates the Lorentz force contribution, this implies $\varepsilon^{\alpha-1}\delta^2 - \delta - \varepsilon^{\alpha+\beta} \gg 0$, which leads to a contradiction as $\delta \rightarrow 0$. On the contrary, assuming that the Lorentz force contribution dominates the free-streaming contribution, this implies $\varepsilon^{\alpha-1}\delta^2 - \delta - \varepsilon^{\alpha+\beta} \ll 0$, which leads also to a contradiction as first $\delta \rightarrow 0$ and next $\varepsilon \rightarrow 0$. In conclusion, if $\alpha\beta + \beta + 3\alpha - 1 > 0$, then the right-hand side of (70) vanishes as $(\varepsilon, \delta) \rightarrow 0$, and we obtain the renormalized Vlasov equation (26).

We continue with the local-in-space entropy conservation law (28). For this purpose, we first restrict entropy functions \mathcal{H} to the set \mathcal{E} , defined by (20), and secondly we take in (70) a test function Φ such that

$$\Phi(t, x, \xi) = \Lambda(t, x)\Theta(\xi) \quad \text{with } \Lambda \in \mathcal{D}((0, T) \times \mathbb{R}^3) \text{ and } \Theta \in \mathcal{D}(\mathbb{R}^3).$$

We then choose the test function Θ such that

$$\Theta(\xi) = \Theta_R(\xi) := \theta(\xi/R) \quad \text{with } R > 0.$$

Here the function $\theta \in \mathcal{D}(\mathbb{R}^3)$ is such that $\text{supp}(\theta) \subset B_{\mathbb{R}^3}(0, 2)$, $\theta \equiv 1$ on $B_{\mathbb{R}^3}(0, 1)$ and $0 \leq \theta \leq 1$ on $B_{\mathbb{R}^3}(0, 2) \setminus B_{\mathbb{R}^3}(0, 1)$. We then have

$$\Theta_R \rightarrow 1, \quad \text{a.e. as } R \rightarrow +\infty \quad \text{and} \quad \nabla_\xi \Theta_R \rightarrow 0, \quad \text{a.e. as } R \rightarrow +\infty. \quad (71)$$

From the uniform integrability assumption (27), and the de La Vallée Poussin theorem, there exists a constant $C_{\mathcal{H}} > 0$, independent of (ε, δ) , but depending on \mathcal{H} such that

$$\int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \leq C_{\mathcal{H}} < +\infty \quad \forall \mathcal{H} \in \mathcal{E}. \quad (72)$$

Using estimate (72), regularity assumptions (23)-(25), and property (71), we obtain from the Lebesgue dominated convergence theorem that

$$\int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \partial_t \Lambda \Theta_R \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \partial_t \Lambda, \quad \text{as } R \rightarrow +\infty, \quad (73)$$

$$\begin{aligned} \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) v^\delta \cdot \nabla_x \Lambda \Theta_R \\ \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) v^\delta \cdot \nabla_x \Lambda, \quad \text{as } R \rightarrow +\infty, \end{aligned} \quad (74)$$

and

$$\mathcal{R}_1 := \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) F_L^{\eta, \varepsilon, \delta} f^{\eta, \varepsilon, \delta} \cdot \nabla_\xi \Theta_R \Lambda \rightarrow 0 \quad \text{as } R \rightarrow +\infty. \quad (75)$$

Limits (73)-(75) are uniform in $(\eta, \varepsilon, \delta)$, and in addition there exists a constant $\kappa_1 > 0$, independent of $(\eta, \varepsilon, \delta)$, but depending on $\|f\|_{L^\infty(0,T;L^2 \cap L^\infty(\mathbb{R}^6))}$, $\|B\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}$, $\|E\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}$, Λ and θ such that

$$|\mathcal{R}_1| \leq \kappa_1 R^{-1}. \quad (76)$$

Using (73)-(76), we obtain from (70),

$$\left| \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) (\partial_t \Lambda + v^\delta \cdot \nabla_x \Lambda) \right| \leq C_* \eta + \kappa_1 R^{-1}. \quad (77)$$

Under the condition, $\alpha\beta + \beta + 3\alpha - 1 > 0$, the right-hand side of (77) vanishes as $(\eta, \varepsilon, \delta) \rightarrow 0$ and $R \rightarrow +\infty$, and we obtain from (77) the local-in-space conservation law (28). In a similar way, by interchanging the role of the test functions Λ and Θ , we obtain local-in-momentum conservation law (29).

We pursue with global entropy conservation law (30). For this aim, we first take in (77) a test function Λ such that

$$\Lambda(t, x) = \varphi(t) \Lambda(x) \quad \text{with } \varphi \in \mathcal{D}((0, T)) \text{ and } \Lambda \in \mathcal{D}(\mathbb{R}^3).$$

We then choose the test function Λ such that

$$\Lambda(x) = \Lambda_R(x) := \lambda(x/R) \quad \text{with } R > 0.$$

Here the function $\lambda \in \mathcal{D}(\mathbb{R}^3)$ is such that $\text{supp}(\lambda) \subset B_{\mathbb{R}^3}(0, 2)$, $\lambda \equiv 1$ on $B_{\mathbb{R}^3}(0, 1)$, and $0 \leq \lambda \leq 1$ on $(B_{\mathbb{R}^3}(0, 2) \setminus B_{\mathbb{R}^3}(0, 1))$. We then have

$$\Lambda_R \rightarrow 1, \quad \text{a.e. as } R \rightarrow +\infty \text{ and } \nabla_x \Lambda_R \rightarrow 0, \quad \text{a.e. as } R \rightarrow +\infty. \quad (78)$$

Using estimate (72), regularity assumptions (23)-(25), and property (78), we obtain from the Lebesgue dominated convergence theorem that

$$\int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \partial_t \varphi \Lambda_R \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \partial_t \varphi \quad \text{as } R \rightarrow +\infty, \quad (79)$$

$$\mathcal{R}_2 := \int_0^T dt \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) v^\delta \cdot \nabla_x \Lambda_R \varphi \rightarrow 0 \quad \text{as } R \rightarrow +\infty. \quad (80)$$

Limits (79)-(80) are uniform in $(\eta, \varepsilon, \delta)$, and in addition there exists a constant $\kappa_2 > 0$, independent of $(\eta, \varepsilon, \delta)$, but depending on $C_{\mathcal{H}}$, φ , and λ such that

$$|\mathcal{R}_2| \leq \kappa_2 R^{-1}. \quad (81)$$

Using (79)-(81), we obtain from (77),

$$\left| \int_0^T dt \partial_t \varphi \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} d\xi \mathcal{H}(f^{\eta, \varepsilon, \delta}) \right| \leq C_* \eta + \kappa_1 R^{-1} + \kappa_2 R^{-1}. \quad (82)$$

Under the condition, $\alpha\beta + \beta + 3\alpha - 1 > 0$, the right-hand side of (82) vanishes as $(\eta, \varepsilon, \delta) \rightarrow 0$ and $R \rightarrow +\infty$, and we obtain from (82) the global entropy conservation law (30). This ends the proof of Theorem 2. \square

4. Energy conservation. As concerns conservation of total energy we have

THEOREM 3. Let (f, E, B) be a weak solution to the relativistic Vlasov–Maxwell system (11)–(14), given by Theorem 1. If the macroscopic kinetic energy density satisfies the supplementary integrability condition,

$$\int_{\mathbb{R}^3} \gamma f d\xi \in L^\infty(0, T; L^2(\mathbb{R}^3)), \quad (83)$$

then, using definition (22), we have the local conservation law of total energy,

$$\partial_t \mathcal{E} + \nabla \cdot \left(\int_{\mathbb{R}^3} \gamma f v d\xi + E \times B \right) = 0 \quad \text{the following in } \mathcal{D}'((0, T) \times \mathbb{R}^3), \quad (84)$$

and the global conservation law of total energy,

$$\mathcal{E}(t) = \mathcal{E}(s) \quad \text{for } 0 < s \leq t < T. \quad (85)$$

REMARK 10. Under assumption (83), it has been proved in [22] that the electromagnetic field (E, B) belongs to $H_{\text{loc}}^s(\mathbb{R}_*^+ \times \mathbb{R}^3)$ with $s = 6/(13 + \sqrt{142})$. Then, such solutions satisfy the conservation laws (84)–(85).

Proof. Choosing in the weak formulation (65) the test function,

$$\Psi(t, x, \xi) = \Lambda(t, x)\Theta(\xi)\gamma(\xi) \in \mathcal{D}((0, T) \times \mathbb{R}^6) \quad \text{with } \Lambda \in \mathcal{D}((0, T) \times \mathbb{R}^3) \text{ and } \Theta \in \mathcal{D}(\mathbb{R}^3), \quad (86)$$

and using $\nabla_\xi \gamma = v$, we obtain

$$\begin{aligned} & \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f \Theta \right) \partial_t \Lambda + \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f v \Theta \right) \cdot \nabla_x \Lambda \\ & + \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi f v \cdot E \Theta \right) \Lambda + \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f F_L \cdot \nabla_\xi \Theta \right) \Lambda = 0. \end{aligned} \quad (87)$$

We now establish the local conservation law of total energy. For this we take in (87) a test function Θ such that

$$\Theta(\xi) = \Theta_R(\xi) := \theta(\xi/R) \quad \text{with } R > 0.$$

Here the function $\theta \in \mathcal{D}(\mathbb{R}^3)$ is such that $\text{supp}(\theta) \subset B_{\mathbb{R}^3}(0, 2)$, $\theta \equiv 1$ on $B_{\mathbb{R}^3}(0, 1)$, and $0 \leq \theta \leq 1$ on $B_{\mathbb{R}^3}(0, 2) \setminus B_{\mathbb{R}^3}(0, 1)$. We then have

$$\Theta_R \rightarrow 1, \quad \text{a.e. as } R \rightarrow +\infty \quad \text{and} \quad \nabla_\xi \Theta_R \rightarrow 0, \quad \text{a.e. as } R \rightarrow +\infty. \quad (88)$$

Using (88) and regularity properties (21), we obtain from the Lebesgue dominated convergence theorem,

$$\int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f \Theta_R \right) \partial_t \Lambda \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f \right) \partial_t \Lambda \quad \text{as } R \rightarrow \infty \quad (89)$$

and

$$\int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f v \Theta_R \right) \cdot \nabla_x \Lambda \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f v \right) \cdot \nabla_x \Lambda \quad \text{as } R \rightarrow \infty. \quad (90)$$

Using assumption (83), regularity properties (21), and Hölder inequality, we obtain

$$\begin{aligned} \left| \int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi \gamma f F_L \cdot \nabla_\xi \Theta_R \right) \Lambda \right| &\leq CR^{-1} \|\nabla \theta\|_{L^\infty} \|\Lambda\|_{L^\infty} \\ \left\| \int_{\mathbb{R}^3} d\xi \gamma f \right\|_{L^\infty(0,T;L^{r'}(\mathbb{R}^3))} (\|E\|_{L^\infty(0,T;L^r(\mathbb{R}^3))} + \|B\|_{L^\infty(0,T;L^r(\mathbb{R}^3))}) &\longrightarrow 0 \quad \text{as } R \rightarrow \infty \end{aligned} \quad (91)$$

with $1/r + 1/r' = 1$ and setting $r = 2$. We now claim that

$$|fv \cdot E\Lambda| \leq |\Lambda| |E| f \in L^\infty(0,T;L^1(\mathbb{R}^6)) \quad \text{if} \quad \int_{\mathbb{R}^3} d\xi \gamma f \in L^\infty(0,T;L^{3/2}(\mathbb{R}^3)). \quad (92)$$

Indeed using interpolation Lemma 2.3 in [22], we obtain

$$\left\| \int_{\mathbb{R}^3} d\xi f \right\|_{L^\infty(0,T;L^2(\mathbb{R}^3))} \leq 9 \|f\|_{L^\infty}^{1/4} \left\| \int_{\mathbb{R}^3} d\xi \gamma f \right\|_{L^\infty(0,T;L^{3/2}(\mathbb{R}^3))}^{3/4}. \quad (93)$$

Therefore, (92) results from (93) and Cauchy-Schwarz inequality. We notice that the $L^{3/2}$ -integrability condition in (92) results from regularity properties (21), assumption (83), and standard interpolation results between Lebesgue spaces. Using (88) we have $fv \cdot E\Theta_R \Lambda \rightarrow fv \cdot E\Lambda$ a.e. as $R \rightarrow +\infty$. Moreover, using (92), we obtain from the Lebesgue dominated convergence theorem,

$$\int_0^T dt \int_{\mathbb{R}^3} dx \left(\int_{\mathbb{R}^3} d\xi fv \cdot E\Theta_R \right) \Lambda \longrightarrow \int_0^T dt \int_{\mathbb{R}^3} dx j \cdot E\Lambda \quad \text{as } R \rightarrow \infty. \quad (94)$$

Using the weak formulation of the Maxwell equation, we obtain

$$\int_0^T dt \int_{\mathbb{R}^3} dx j \cdot E\Lambda = \int_0^T dt \int_{\mathbb{R}^3} dx \frac{|E|^2 + |B|^2}{2} \partial_t \Lambda + \int_0^T dt \int_{\mathbb{R}^3} dx E \times B \cdot \nabla_x \Lambda. \quad (95)$$

Using (89)-(91) and (94)-(95), we obtain from (87),

$$\begin{aligned} \int_0^T dt \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma f \right) + \frac{|E|^2 + |B|^2}{2} \right\} \partial_t \Lambda \\ + \int_0^T dt \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma fv \right) + E \times B \right\} \cdot \nabla_x \Lambda = 0, \end{aligned} \quad (96)$$

which gives the local conservation law of total energy (84). We continue by deriving the global conservation law of total energy. For this we take in (96) a test function Λ such that

$$\Lambda(t, x) = \varphi(t)\Lambda(x) \quad \text{with } \varphi \in \mathcal{D}((0, T)) \quad \text{and} \quad \Lambda \in \mathcal{D}(\mathbb{R}^3).$$

We then choose the test function Λ such that

$$\Lambda(x) = \Lambda_R(x) := \lambda(x/R) \quad \text{with } R > 0.$$

Here the function $\lambda \in \mathcal{D}(\mathbb{R}^3)$ is such that $\text{supp}(\lambda) \subset B_{\mathbb{R}^3}(0, 2)$, $\lambda \equiv 1$ on $B_{\mathbb{R}^3}(0, 1)$, and $0 \leq \lambda \leq 1$ on $(B_{\mathbb{R}^3}(0, 2) \setminus B_{\mathbb{R}^3}(0, 1))$. We then have

$$\Lambda_R \longrightarrow 1, \quad \text{a.e. as } R \rightarrow +\infty \quad \text{and} \quad \nabla_x \Lambda_R \longrightarrow 0, \quad \text{a.e. as } R \rightarrow +\infty. \quad (97)$$

Using (97) and regularity properties (21), especially $f \in L^\infty(0, T; L_\gamma^1(\mathbb{R}^6))$ and $E, B \in L^\infty(0, T; L^2(\mathbb{R}^3))$, we obtain from the Lebesgue dominated convergence theorem,

$$\begin{aligned} & \int_0^T dt \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma f \right) + \frac{|E|^2 + |B|^2}{2} \right\} \Lambda_R \partial_t \varphi \\ & \quad \rightarrow \int_0^T dt \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma f \right) + \frac{|E|^2 + |B|^2}{2} \right\} \partial_t \varphi \quad \text{as } R \rightarrow +\infty \end{aligned} \quad (98)$$

and

$$\int_0^T dt \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma f v \right) + E \times B \right\} \cdot \nabla_x \Lambda_R \varphi \rightarrow 0 \quad \text{as } R \rightarrow +\infty. \quad (99)$$

Using (98)-(99), and passing to the limit $R \rightarrow +\infty$ in (96), with $\Lambda(t, x) = \varphi(t) \lambda(x/R)$, we obtain

$$\int_0^T dt \partial_t \varphi \int_{\mathbb{R}^3} dx \left\{ \left(\int_{\mathbb{R}^3} d\xi \gamma f \right) + \frac{|E|^2 + |B|^2}{2} \right\} = 0, \quad (100)$$

which gives the global conservation law of total energy (85). \square

REMARK 11. 1. If $E, B \in L^\infty(0, T; L^\infty(\mathbb{R}^6))$, we observe that the proof of Theorem 3 remains valid without condition (83), and then local and global conservation of total energy (84)-(85) are satisfied.

2. Using the continuous embedding $W^{\beta, q}(\mathbb{R}^3) \subset L^{3q/(3-\beta q)}(\mathbb{R}^3)$, with $\beta q < 3$, we observe that if $E, B \in L^\infty(0, T; L^2 \cap W^{\beta, q}(\mathbb{R}^3))$, and

$$\int_{\mathbb{R}^3} d\xi \gamma f \in L^\infty(0, T; L^{3q/((3+\beta)q-3)}(\mathbb{R}^3)),$$

then estimates (91) and (94) still hold. Therefore, local and global conservation of total energy (84)-(85) are satisfied.

Acknowledgments. The authors would like to thank Gregory Eyink for his constructive comments on this work. The first and third authors wish to thank the Observatoire de la Côte d'Azur and the Laboratoire J.-L. Lagrange for their hospitality and financial support. The third author's research was supported by the NSF under grant DMS-1764119 and by an AMS Centennial Fellowship. Part of this work was done while the third author was visiting the Department of Mathematics and the Program in Applied and Computational Mathematics at Princeton University.

REFERENCES

- [1] R. A. Adams, *Sobolev spaces*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65. MR0450957
- [2] M. Aizenman, *On vector fields as generators of flows: a counterexample to Nelson's conjecture*, Ann. Math. (2) **107** (1978), no. 2, 287–296, DOI 10.2307/1971145. MR0482853
- [3] I. Akramov and E. Wiedemann, *Renormalization of active scalar equations*, Nonlinear Anal. **179** (2019), 254–269, DOI 10.1016/j.na.2018.08.018. MR3886632
- [4] G. Alberti, G. Crippa, and A. L. Mazzucato, *Exponential self-similar mixing and loss of regularity for continuity equations* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **352** (2014), no. 11, 901–906, DOI 10.1016/j.crma.2014.08.021. MR3268760
- [5] G. Alberti, G. Crippa, and A. L. Mazzucato, *Exponential self-similar mixing by incompressible flows*, J. Amer. Math. Soc. **32** (2019), no. 2, 445–490, DOI 10.1090/jams/913. MR3904158

- [6] G. Alberti, S. Bianchini, and G. Crippa, *Structure of level sets and Sard-type properties of Lipschitz maps*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **12** (2013), no. 4, 863–902. MR3184572
- [7] G. Alberti, S. Bianchini, and G. Crippa, *A uniqueness result for the continuity equation in two dimensions*, J. Eur. Math. Soc. (JEMS) **16** (2014), no. 2, 201–234, DOI 10.4171/JEMS/431. MR3161282
- [8] S. Alinhac and P. Gérard, *Opérateurs pseudo-différentiels et théorème de Nash-Moser* (French), Savoirs Actuels. [Current Scholarship], InterEditions, Paris; Éditions du Centre National de la Recherche Scientifique (CNRS), Meudon, 1991. MR1172111
- [9] L. Ambrosio, *Transport equation and Cauchy problem for BV vector fields*, Invent. Math. **158** (2004), no. 2, 227–260, DOI 10.1007/s00222-004-0367-2. MR2096794
- [10] L. Ambrosio, *Well posedness of ODE's and continuity equations with nonsmooth vector fields, and applications*, Rev. Mat. Complut. **30** (2017), no. 3, 427–450, DOI 10.1007/s13163-017-0244-3. MR3690748
- [11] L. Ambrosio, M. Colombo, and A. Figalli, *Existence and uniqueness of maximal regular flows for non-smooth vector fields*, Arch. Ration. Mech. Anal. **218** (2015), no. 2, 1043–1081, DOI 10.1007/s00205-015-0875-9. MR3375545
- [12] L. Ambrosio, M. Colombo, and A. Figalli, *On the Lagrangian structure of transport equations: the Vlasov-Poisson system*, Duke Math. J. **166** (2017), no. 18, 3505–3568, DOI 10.1215/00127094-2017-0032. MR3732882
- [13] L. Ambrosio and G. Crippa, *Continuity equations and ODE flows with non-smooth velocity*, Proc. Roy. Soc. Edinburgh Sect. A **144** (2014), no. 6, 1191–1244, DOI 10.1017/S0308210513000085. MR3283066
- [14] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace transforms and Cauchy problems*, 2nd ed., Monographs in Mathematics, vol. 96, Birkhäuser/Springer Basel AG, Basel, 2011. MR2798103
- [15] J.-M. Aubry, D. Maman, and S. Seuret, *Local behavior of traces of Besov functions: prevalent results*, J. Funct. Anal. **264** (2013), no. 3, 631–660, DOI 10.1016/j.jfa.2012.11.012. MR3003731
- [16] C. Bardos and E. S. Titi, *Loss of smoothness and energy conserving rough weak solutions for the 3d Euler equations*, Discrete Contin. Dyn. Syst. Ser. S **3** (2010), no. 2, 185–197, DOI 10.3934/dcdss.2010.3.185. MR2610558
- [17] C. Bardos, E. S. Titi, and E. Wiedemann, *The vanishing viscosity as a selection principle for the Euler equations: the case of 3D shear flow* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **350** (2012), no. 15–16, 757–760, DOI 10.1016/j.crma.2012.09.005. MR2981348
- [18] C. Bardos and E. S. Titi, *Onsager's conjecture for the incompressible Euler equations in bounded domains*, Arch. Ration. Mech. Anal. **228** (2018), no. 1, 197–207, DOI 10.1007/s00205-017-1189-x. MR3749259
- [19] C. Bardos, E. S. Titi, and E. Wiedemann, *Onsager's conjecture with physical boundaries and an application to the vanishing viscosity limit*, Preprint, 2018, arXiv:1803.04939.
- [20] C. Bardos, P. Gwiazda, A. Świerniewska-Gwiazda, E. S. Titi, and E. Wiedemann, *On the extension of Onsager's conjecture for general conservation laws*, J. Nonlinear Sci. **29** (2019), no. 2, 501–510, DOI 10.1007/s00332-018-9496-4. MR3927104
- [21] J. Bergh and J. Löfström, *Interpolationsprostora (Russian)*, “Mir”, Moscow, 1980. Vvedenie. [An introduction]; Translated from the English by V. S. Krjučkov and P. I. Lizorkin. MR608963
- [22] N. Besse and P. Bechouche, *Regularity of weak solutions for the relativistic Vlasov-Maxwell system*, J. Hyperbolic Differ. Equ. **15** (2018), no. 4, 693–719, DOI 10.1142/S0219891618500212. MR3900881
- [23] F. Bouchut, *Renormalized solutions to the Vlasov equation with coefficients of bounded variation*, Arch. Ration. Mech. Anal. **157** (2001), no. 1, 75–90, DOI 10.1007/PL00004237. MR1822415
- [24] F. Bouchut, F. Golse, and M. Pulvirenti, *Kinetic equations and asymptotic theory*, Series in Applied Mathematics (Paris), vol. 4, Gauthier-Villars, Éditions Scientifiques et Médicales Elsevier, Paris, 2000. Edited and with a foreword by Benoît Perthame and Laurent Desvillettes. MR2065070
- [25] F. Bouchut, F. Golse, and C. Pallard, *Nonresonant smoothing for coupled wave + transport equations and the Vlasov-Maxwell system*, Rev. Mat. Iberoamericana **20** (2004), no. 3, 865–892, DOI 10.4171/RMI/409. MR2124491
- [26] J. Brasseur, *On restrictions of Besov functions*, Nonlinear Anal. **170** (2018), 197–225, DOI 10.1016/j.na.2018.01.004. MR3765561

- [27] A. Cheskidov, P. Constantin, S. Friedlander, and R. Shvydkoy, *Energy conservation and Onsager's conjecture for the Euler equations*, Nonlinearity **21** (2008), no. 6, 1233–1252, DOI 10.1088/0951-7715/21/6/005. MR2422377
- [28] F. Colombini and N. Lerner, *Uniqueness of continuous solutions for BV vector fields*, Duke Math. J. **111** (2002), no. 2, 357–384, DOI 10.1215/S0012-7094-01-11126-5. MR1882138
- [29] F. Colombini, T. Luo, and J. Rauch, *Uniqueness and nonuniqueness for nonsmooth divergence free transport*, Séminaire: Équations aux Dérivées Partielles, 2002–2003, Sémin. Équ. Dériv. Partielles, École Polytech., Palaiseau, 2003, pp. Exp. No. XXII, 21. MR2030717
- [30] G. Crippa, N. Gusev, S. Spirito, and E. Wiedemann, *Non-uniqueness and prescribed energy for the continuity equation*, Commun. Math. Sci. **13** (2015), no. 7, 1937–1947, DOI 10.4310/CMS.2015.v13.n7.a12. MR3393180
- [31] G. Crippa, N. Gusev, S. Spirito, and E. Wiedemann, *Failure of the chain rule for the divergence of bounded vector fields*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **17** (2017), no. 1, 1–18. MR3676038
- [32] P. Constantin, W. E, and E. S. Titi, *Onsager's conjecture on the energy conservation for solutions of Euler's equation*, Comm. Math. Phys. **165** (1994), no. 1, 207–209. MR1298949
- [33] N. Depauw, *Non unicité des solutions bornées pour un champ de vecteurs BV en dehors d'un hyperplan* (French, with English and French summaries), C. R. Math. Acad. Sci. Paris **337** (2003), no. 4, 249–252, DOI 10.1016/S1631-073X(03)00330-3. MR2009116
- [34] R. DiPerna and P.-L. Lions, *Solutions globales d'équations du type Vlasov-Poisson* (French, with English summary), C. R. Acad. Sci. Paris Sér. I Math. **307** (1988), no. 12, 655–658. MR967806
- [35] R. J. DiPerna and P.-L. Lions, *Global weak solutions of kinetic equations*, Rend. Sem. Mat. Univ. Politec. Torino **46** (1988), no. 3, 259–288 (1990). MR1101105
- [36] R. J. DiPerna and P.-L. Lions, *Global weak solutions of Vlasov-Maxwell systems*, Comm. Pure Appl. Math. **42** (1989), no. 6, 729–757, DOI 10.1002/cpa.3160420603. MR1003433
- [37] R. J. DiPerna and P.-L. Lions, *Ordinary differential equations, transport theory and Sobolev spaces*, Invent. Math. **98** (1989), no. 3, 511–547, DOI 10.1007/BF01393835. MR1022305
- [38] R. J. DiPerna and P.-L. Lions, *On the Cauchy problem for Boltzmann equations: global existence and weak stability*, Ann. of Math. (2) **130** (1989), no. 2, 321–366, DOI 10.2307/1971423. MR1014927
- [39] T. D. Drivas and G. L. Eyink, *An Onsager singularity theorem for turbulent solutions of compressible Euler equations*, Comm. Math. Phys. **359** (2018), no. 2, 733–763, DOI 10.1007/s00220-017-3078-4. MR3783560
- [40] T. D. Drivas and H. Q. Nguyen, *Onsager's conjecture and anomalous dissipation on domains with boundary*, SIAM J. Math. Anal. **50** (2018), no. 5, 4785–4811, DOI 10.1137/18M1178864. MR3851045
- [41] J. Duchon and R. Robert, *Inertial energy dissipation for weak solutions of incompressible Euler and Navier-Stokes equations*, Nonlinearity **13** (2000), no. 1, 249–255, DOI 10.1088/0951-7715/13/1/312. MR1734632
- [42] K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. MR1721989
- [43] G. L. Eyink, *Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer*, Phys. D **78** (1994), no. 3-4, 222–240, DOI 10.1016/0167-2789(94)90117-1. MR1302409
- [44] G. L. Eyink, *Cascades and dissipative anomalies in nearly Collisionless Plasma turbulence*, Phys. Rev. X **8** (2018) 041020.
- [45] E. Feireisl, P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann, *Regularity and energy conservation for the compressible Euler equations*, Arch. Ration. Mech. Anal. **223** (2017), no. 3, 1375–1395, DOI 10.1007/s00205-016-1060-5. MR3594358
- [46] X. Fernández-Real, *The Lagrangian structure of the Vlasov-Poisson system in domains with specular reflection*, Comm. Math. Phys. **364** (2018), no. 3, 1327–1406, DOI 10.1007/s00220-018-3242-5. MR3875828
- [47] U. S. Fjordholm and E. Wiedemann, *Statistical solutions and Onsager's conjecture*, Phys. D **376/377** (2018), 259–265, DOI 10.1016/j.physd.2017.10.009. MR3815222
- [48] R. T. Glassey, *The Cauchy problem in kinetic theory*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1996. MR1379589
- [49] Y. Guo, *Global weak solutions of the Vlasov-Maxwell system with boundary conditions*, Comm. Math. Phys. **154** (1993), no. 2, 245–263. MR1224079

- [50] Y. Guo, *Regularity for the Vlasov equations in a half-space*, Indiana Univ. Math. J. **43** (1994), no. 1, 255–320, DOI 10.1512/iumj.1994.43.43013. MR1275462
- [51] Y. Guo, *Singular solutions of the Vlasov-Maxwell system on a half line*, Arch. Rational Mech. Anal. **131** (1995), no. 3, 241–304, DOI 10.1007/BF00382888. MR1354697
- [52] P. Gwiazda, M. Michálek, and A. Świernczewska-Gwiazda, *A note on weak solutions of conservation laws and energy/entropy conservation*, Arch. Ration. Mech. Anal. **229** (2018), no. 3, 1223–1238, DOI 10.1007/s00205-018-1238-0. MR3814601
- [53] M. Hauray, *On Liouville transport equation with force field in BV_{loc}* , Comm. Partial Differential Equations **29** (2004), no. 1-2, 207–217, DOI 10.1081/PDE-120028850. MR2038150
- [54] S. Jaffard, *Théorèmes de trace et “dimensions négatives”* (French, with English and French summaries), C. R. Acad. Sci. Paris Sér. I Math. **320** (1995), no. 4, 409–413. MR1320112
- [55] C. Le Bris and P.-L. Lions, *Renormalized solutions of some transport equations with partially $W^{1,1}$ velocities and applications*, Ann. Mat. Pura Appl. (4) **183** (2004), no. 1, 97–130, DOI 10.1007/s10231-003-0082-4. MR2044334
- [56] T. M. Leslie and R. Shvydkoy, *The energy balance relation for weak solutions of the density-dependent Navier-Stokes equations*, J. Differential Equations **261** (2016), no. 6, 3719–3733, DOI 10.1016/j.jde.2016.06.001. MR3527643
- [57] P.-L. Lions, *Mathematical topics in fluid mechanics. Vol. 2*, Oxford Lecture Series in Mathematics and its Applications, vol. 10, The Clarendon Press, Oxford University Press, New York, 1998. Compressible models; Oxford Science Publications. MR1637634
- [58] L. Onsager, *Statistical hydrodynamics*, Nuovo Cimento (9) **6** (1949), no. Supplemento, 2 (Convegno Internazionale di Meccanica Statistica), 279–287. MR0036116
- [59] G. Rein, *Global weak solutions to the relativistic Vlasov-Maxwell system revisited*, Commun. Math. Sci. **2** (2004), no. 2, 145–158. MR2119935
- [60] J. C. Robinson, J. L. Rodrigo, and J. W. D. Skipper, *Energy conservation in the 3D Euler equation on $\mathbb{T}^2 \times \mathbb{R}_+$* , Partial differential equations in fluid mechanics, London Math. Soc. Lecture Note Ser., vol. 452, Cambridge Univ. Press, Cambridge, 2018, pp. 224–251. MR3838052
- [61] J. C. Robinson, J. L. Rodrigo, and J. W. D. Skipper, *Energy conservation for the Euler equations on $\mathbb{T}^2 \times \mathbb{R}_+$ for weak solutions defined without reference to the pressure*, Asymptot. Anal. **110** (2018), no. 3-4, 185–202, DOI 10.3233/ASY-181482. MR3885495
- [62] L. Székelyhidi, *Weak solutions to the incompressible Euler equations with vortex sheet initial data* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **349** (2011), no. 19–20, 1063–1066, DOI 10.1016/j.crma.2011.09.009. MR2842999
- [63] H. Triebel, *Theory of function spaces*, Monographs in Mathematics, vol. 78, Birkhäuser Verlag, Basel, 1983. MR781540
- [64] H. Triebel, *The structure of functions*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2001. [2012 reprint of the 2001 original] [MR1851996]. MR3013187
- [65] Y. Yao and A. Zlatoš, *Mixing and un-mixing by incompressible flows*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 7, 1911–1948, DOI 10.4171/JEMS/709. MR3656475
- [66] K. Yosida, *Functional analysis*, 6th ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 123, Springer-Verlag, Berlin-New York, 1980. MR617913
- [67] C. Yu, *Energy conservation for the weak solutions of the compressible Navier-Stokes equations*, Arch. Ration. Mech. Anal. **225** (2017), no. 3, 1073–1087, DOI 10.1007/s00205-017-1121-4. MR3667283